

Resonant inelastic scattering study of insulating cuprates at the Cu L-edges

M.-Z. Hasan¹, E. D. Isaacs²

A. Agui³, C. S  the³, T. K   mbre³, J. Rubensson³, J. Nordgren³

E. Gullickson⁴, P. Platzman², Z.-X. Shen¹

L. L. Miller⁵, S.W. Cheong²

¹Dept. of Applied Physics, Stanford University, Stanford, CA 94305

²Lucent Technologies, Bell Laboratories, Murray Hill, NJ 07975

⁴Lawrence Berkeley National Laboratory, Berkeley, CA 94720

³Dept. of Physics, Uppsala Univ., Box 530, S-75121 Uppsala, Sweden

⁵Ames Laboratory, Ames, IA 50011

We have performed some initial measurements of resonant inelastic scattering spectra for three parent cuprates (La_2CuO_4 , $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ and $\text{Ca}_2\text{CuO}_2\text{Cl}_2$) at the Cu L2 & L3 edges. They all show similar features - a very strong excitation band of dd type around 2 eV and a charge-transfer excitation feature around 5 eV. L-edge spectra have larger intensity at the dd channel compared to the Cu K-edge results we have from Brookhaven [1,2]. Such resonant profile measurements are important to characterize the dd excitations channels (optically forbidden) in these compounds as they are relevant for antiferromagnetic and superconducting properties which arise when these insulators are doped with different concentrations. More detailed study is underway.

[1] P. Abbamonte, E.D. Isaacs et.al. (submitted to PRL)

[2] Z. Hasan & E. D. Issacs (to be published)

This work was supported by Lucent Technologies, Bell Laboratories.

Principal investigator: Eric D. Isaacs, Lucent Technologies, Bell Laboratories. Email: edi@physics.bell-labs.com. Telephone: 908-582-7261.